

On the γ -reflected Processes with fBm Input

Peng Liu*, Enkelejd Hashorva[†] and Lanpeng Ji[†]

February 12, 2014

Abstract: Define a γ -reflected process $W_\gamma(t) = Y_H(t) - \gamma \inf_{s \in [0, t]} Y_H(s)$, $t \geq 0$ with input process $\{Y_H(t), t \geq 0\}$ which is a fractional Brownian motion with Hurst index $H \in (0, 1)$ and a negative linear trend. In risk theory $R_\gamma(t) = u - W_\gamma(t)$, $t \geq 0$ is referred to as the risk process with tax of a loss-carry-forward type, whereas in queueing theory W_1 is referred to as the queue length process. In this paper, we investigate the ruin probability and the ruin time of the risk process R_γ over a surplus dependent time interval $[0, T_u]$.

Key Words: γ -reflected process; risk process with tax; ruin time; maximum losses; fractional Brownian motion; Piterbarg constant; Piterbarg's theorem; Pickands constant.

AMS Classification: Primary 60G15; secondary 60G70

1 Introduction

Let $\{X_H(t), t \geq 0\}$ be a standard fractional Brownian motion (fBm) with Hurst index $H \in (0, 1)$ meaning that X_H is a centered Gaussian process with almost surely continuous sample paths and covariance function

$$Cov(X_H(t), X_H(s)) = \frac{1}{2}(|t|^{2H} + |s|^{2H} - |t - s|^{2H}), \quad t, s \geq 0$$

We define a γ -reflected process with input process $Y_H(t) = X_H(t) - ct$, $c > 0$ by

$$W_\gamma(t) = Y_H(t) - \gamma \inf_{s \in [0, t]} Y_H(s), \quad t \geq 0, \quad (1)$$

where $\gamma \in [0, 1]$ is the reflection parameter.

Motivations for studying W_γ come from its wide applications in the fields of queueing, insurance,

*School Mathematical Sciences and LMPC, Nankai University, Tianjin 300071, China

[†]Department of Actuarial Science, University of Lausanne, UNIL-Dorigny 1015 Lausanne, Switzerland

finance and telecommunication. For instance, in queuing theory W_1 is referred to as the queue length process (or the workload process); see, e.g., [17, 2, 26, 25, 4] among many others. In risk theory the process $R_\gamma(t) = u - W_\gamma(t), t \geq 0, u \geq 0$ is referred to as the risk process with tax payments of a loss-carry-forward type; see, e.g., [3]. We refer to [10, 11, 20, 21, 22, 12] for some recent studies of W_0 .

For any $u \geq 0$, define the *ruin time* of the γ -reflected process W_γ by

$$\tau_{\gamma,u} = \inf\{t \geq 0 : W_\gamma(t) > u\} \quad (\text{with } \inf\{\emptyset\} = \infty). \quad (2)$$

Further let $T_u, u \geq 0$ be a positive function and define the *ruin probability* over a surplus dependent time interval $[0, T_u]$ by

$$\psi_{\gamma,T_u}(u) := \mathbb{P}(\tau_{\gamma,u} \leq T_u).$$

Hereafter, $\psi_{\gamma,\infty}(u)$ denotes the ruin probability over an infinite-time horizon.

The ruin time and the ruin probability for the case that $T_u \equiv T \in (0, \infty)$ and the case that $T_u = \infty$ are studied in [19, 18]; see also [11, 20, 21, 22]. In [19] the exact asymptotics of $\psi_{\gamma,T}(u)$ and $\psi_{\gamma,\infty}(u)$ are derived, which combined with the results in [20] and [11] lead to the following asymptotic equivalence

$$\psi_{\gamma,T}(u) = \mathcal{C}_{H,\gamma} \psi_{0,T}(u)(1 + o(1)), \quad u \rightarrow \infty \quad (3)$$

for any $T \in (0, \infty]$, with $\mathcal{C}_{H,\gamma}$ some known positive constant. The recent contribution [18] investigates the approximation of the conditional ruin time $\tau_{\gamma,u} | (\tau_{\gamma,u} < \infty)$. As shown therein the following convergence in distribution (denoted by $\xrightarrow{d}$)

$$\frac{\tau_{\gamma,u} - t_0 u}{A(u)} \Big| (\tau_{\gamma,u} < \infty) \xrightarrow{d} \mathcal{N} \quad (4)$$

holds as $u \rightarrow \infty$ for any $\gamma \in [0, 1)$, where $\mathcal{N}$ is an $N(0, 1)$ random variable and

$$t_0 = \frac{H}{c(1-H)}, \quad A(u) = \frac{H^{H+1/2} u^H}{(1-H)^{H+\frac{1}{2}} c^{H+1}}. \quad (5)$$

See also [21, 22, 23, 8] for related results. We note in passing that the ruin time and the ruin probability are also studied extensively in the framework of other stochastic processes; see, e.g., [15, 16, 14, 3].

With motivation from [5] and [9], as a continuation of the investigation of the aforementioned papers we shall analyze the ruin probability and the conditional ruin time of W_γ over the surplus

dependent time interval $[0, T_u]$ letting $u \rightarrow \infty$. In the literature, commonly the case $T = \infty$ is considered since for many models explicit calculations of the ruin probability is possible. From a practical point of view, a more interesting quantity is the finite-time ruin probability. The case of the surplus dependent time interval lies thus in between and is of both practical and theoretical interests; results for the ruin probability in this case (with $\gamma = 0$) are initially derived in [5], see Theorem 2.1 below.

The novel aspect of this paper is that T_u will be a function changing with u according to three different scenarios which we can define with help of (4). In Theorem 2.2 below we show that similar asymptotic equivalence as in (3) still holds for all the three scenarios. In Theorem 2.3 we derive a truncated Gaussian approximation for the (scaled) conditional ruin time over the long time horizon, whereas for the short and the intermediate time horizons an exponential approximation is possible.

Organization of the rest of the paper: The main results are presented in Section 2 followed then by a section dedicated to the proofs. In Appendix we present a variant of the celebrated Piterbarg's theorem, which is of interest for further theoretical developments.

2 Main Results

In this contribution, three scenarios of T_u will be distinguished. In view of (4) the asymptotic mean of the ruin time (equal to $t_0 u$) and the asymptotic standard deviation $A(u)$ (see (5)) should be used as a scaling parameter for T_u leading to the definition of the following three scenarios:

- i) The short time horizon: $\lim_{u \rightarrow \infty} T_u/u = 0$;
- ii) The intermediate time horizon: $\lim_{u \rightarrow \infty} T_u/u = s_0 \in (0, t_0)$;
- iii) The long time horizon: $\lim_{u \rightarrow \infty} \frac{T_u - t_0 u}{A(u)} = x \in (-\infty, \infty]$.

Next we introduce two well-known constants appearing in the asymptotic theory of Gaussian processes. Let therefore $\{B_\alpha(t), t \geq 0\}$ be a standard fBm with Hurst index $\alpha/2 \in (0, 1]$. The *Pickands constant* is defined by

$$\mathcal{H}_\alpha = \lim_{S \rightarrow \infty} \frac{1}{S} \mathbb{E} \left(\exp \left(\sup_{t \in [0, S]} \left(\sqrt{2} B_\alpha(t) - t^\alpha \right) \right) \right) \in (0, \infty), \quad \alpha \in (0, 2]$$

and the *Piterbarg constant* is given by

$$\mathcal{P}_\alpha^b = \lim_{S \rightarrow \infty} \mathbb{E} \left(\exp \left(\sup_{t \in [0, S]} \left(\sqrt{2} B_\alpha(t) - (1+b)t^\alpha \right) \right) \right) \in (0, \infty), \quad \alpha \in (0, 2], \quad b > 0.$$

We refer to [24, 7, 10, 6, 1, 19, 13] for properties and extensions of the Pickands and Piterbarg constants.

In what follows denote by $\Phi(\cdot)$ the distribution function of an $N(0, 1)$ random variable, write $\mathcal{N}$ for an $N(0, 1)$ random variable and put $\Psi(\cdot) := 1 - \Phi(\cdot)$. Before displaying our main results, we include below a key finding of [5] concerning the ruin probability of the 0-reflected process W_0 .

Theorem 2.1 *Let W_0 be the 0-reflected process given as in (1) with $H \in (0, 1)$. We have*

i) *If $\lim_{u \rightarrow \infty} T_u/u = s_0 \in [0, t_0)$, then*

$$\psi_{0, T_u}(u) = D_H \left(\frac{u + cT_u}{T_u^H} \right)^{\left(\frac{1-2H}{H}\right)_+} \Psi \left(\frac{u + cT_u}{T_u^H} \right) (1 + o(1)) \quad (6)$$

as $u \rightarrow \infty$, where

$$D_H = \begin{cases} 2^{-\frac{1}{2H}} (H - c_0)^{-1} \mathcal{H}_{2H}, & \text{if } H < 1/2, \\ \frac{4(1-c_0)^2}{(1-2c_0)(2-2c_0)}, & \text{if } H = 1/2, \\ 1 & \text{if } H > 1/2, \end{cases} \quad \text{with } c_0 = \frac{cs_0}{1 + cs_0}.$$

ii) *If $\lim_{u \rightarrow \infty} \frac{T_u - t_0 u}{A(u)} = x \in (-\infty, \infty]$, then*

$$\psi_{0, T_u}(u) = \psi_{0, \infty}(u) \Phi(x) (1 + o(1)) \quad (7)$$

as $u \rightarrow \infty$, where the infinite-time ruin probability $\psi_{0, \infty}(u)$ is given by

$$\psi_{0, \infty}(u) = 2^{\frac{1}{2} - \frac{1}{2H}} \frac{\sqrt{\pi}}{\sqrt{H(1-H)}} \mathcal{H}_{2H} \left(\frac{c^H u^{1-H}}{H^H (1-H)^{1-H}} \right)^{1/H-1} \Psi \left(\frac{c^H u^{1-H}}{H^H (1-H)^{1-H}} \right) (1 + o(1)). \quad (8)$$

The next theorem shows the asymptotic relations between the ruin probability of the γ -reflected process W_γ and that of the 0-reflected process W_0 . Therefore in the light of Theorem 2.1 we obtain the exact asymptotics of the ruin probability over the surplus dependent interval $[0, T_u]$ of the γ -reflected process W_γ .

Theorem 2.2 *Let W_γ be the γ -reflected process given as in (1) with $H \in (0, 1)$ and $\gamma \in (0, 1)$.*

We have

i) *If $\lim_{u \rightarrow \infty} T_u/u = s_0 \in [0, t_0)$, then*

$$\psi_{\gamma, T_u}(u) = \mathcal{M}_{H, \gamma} \psi_{0, T_u}(u) (1 + o(1)) \quad (9)$$

as $u \rightarrow \infty$, where

$$\mathcal{M}_{H, \gamma} = \begin{cases} \mathcal{P}_{2H}^{\frac{1-\gamma}{\gamma}}, & \text{if } H < 1/2, \\ \frac{2-2c_0}{2-2c_0-\gamma}, & \text{if } H = 1/2, \\ 1 & \text{if } H > 1/2. \end{cases}$$

ii) If $\lim_{u \rightarrow \infty} \frac{T_u - t_0 u}{A(u)} = x \in (-\infty, \infty]$, then

$$\psi_{\gamma, T_u}(u) = \mathcal{P}_{2H}^{\frac{1-\gamma}{\gamma}} \psi_{0, T_u}(u)(1 + o(1)) \quad (10)$$

as $u \rightarrow \infty$.

Remarks. a) For the case that $\gamma = 1$ we can add: Under the statement i) above similar arguments as in the proof of Theorem 2.2 show that (9) holds as $u \rightarrow \infty$, with

$$\mathcal{M}_{H,1} = \begin{cases} 2^{-\frac{1}{2H}}(H - c_0)^{-1} \mathcal{H}_{2H}, & \text{if } H < 1/2, \\ \frac{2-2c_0}{1-2c_0}, & \text{if } H = 1/2, \\ 1 & \text{if } H > 1/2. \end{cases}$$

For ii), depending on the values of x different asymptotics will appear; those derivations are more involved and will therefore be omitted here.

b) Another scenario of T_u which is between the cases i) and ii) is that $\lim_{u \rightarrow \infty} \frac{T_u - t_0 u}{A(u)} = -\infty$. This case can not be dealt with in general; more conditions should be imposed for the asymptotic behaviour of T_u around $t_0 u$.

c) As discussed in [5] also of interest is the investigation of the maximum losses given that ruin occurs, which, in our setup, is defined as

$$L(\gamma, u) := \left(\sup_{t \in [0, T_u]} W_\gamma(t) - u \right) \Big|_{(\tau_u \leq T_u)}. \quad (11)$$

Under the assumptions of Theorem 2.2, we have by an application of Theorem 2.1 and Theorem 2.2 that if i) is satisfied, then

$$\frac{(u + cT_u)L(\gamma, u)}{T_u^{2H}} \xrightarrow{d} \mathcal{E}, \quad u \rightarrow \infty,$$

and if ii) is valid, then

$$\frac{c^{2H}(1-H)^{2H-1}L(\gamma, u)}{H^{2H}u^{2H-1}} \xrightarrow{d} \mathcal{E}, \quad u \rightarrow \infty.$$

Here (and in the sequel) $\mathcal{E}$ denotes a unit exponential random variable. Note in passing that the last convergence in distribution is clear when $\gamma = 0$ and $T_u = \infty$ since it is known that the random variable $\sup_{t \in [0, \infty)} W_0(t)$ is exponentially distributed with parameter $2c$.

Below we shall establish asymptotic approximations for the ruin times considering all three scenarios for T_u . It turns out that for the long time horizon the (scaled) conditional ruin time can be approximated by a truncated Gaussian random variable. Surprisingly, this is no longer the case for the short and the intermediate time horizons where the (scaled) conditional ruin time is approximated by an exponential random variable.

Theorem 2.3 Let W_γ be the γ -reflected process given as in (1) with $H \in (0, 1)$ and $\gamma \in (0, 1)$, and let $\tau_{\gamma,u}$ be the ruin time defined as in (2). We have

i) If $\lim_{u \rightarrow \infty} T_u/u = 0$, then

$$\frac{Hu^2(T_u - \tau_{\gamma,u})}{T_u^{2H+1}} \Big| (\tau_{\gamma,u} \leq T_u) \xrightarrow{d} \mathcal{E}, \quad u \rightarrow \infty;$$

ii) If $\lim_{u \rightarrow \infty} T_u/u = s_0 \in (0, t_0)$, then

$$\frac{(1 + cs_0)(H - (1 - H)cs_0)(T_u - \tau_{\gamma,u})}{s_0^{2H+1}u^{2H-1}} \Big| (\tau_{\gamma,u} \leq T_u) \xrightarrow{d} \mathcal{E}, \quad u \rightarrow \infty;$$

iii) If $\lim_{u \rightarrow \infty} \frac{T_u - t_0 u}{A(u)} = x \in (-\infty, \infty]$, then

$$\frac{\tau_{\gamma,u} - t_0 u}{A(u)} \Big| (\tau_{\gamma,u} \leq T_u) \xrightarrow{d} \mathcal{N} \Big| (\mathcal{N} < x), \quad u \rightarrow \infty.$$

Remark. As expected, the approximation of the conditional ruin time does not involve the reflection constant γ , since in view of the proof of Theorem 2.3 the terms with γ are canceled out because of the conditional event.

3 Proofs

In this section, we shall present the proofs of all the theorems. We start with the proof of Theorem

2.2. First note that for any $u > 0$

$$\begin{aligned} \psi_{\gamma, T_u}(u) &= \mathbb{P} \left(\sup_{t \in [0, T_u]} W_\gamma(t) > u \right) \\ &= \mathbb{P} \left(\sup_{0 \leq s \leq t \leq T_u} (Z(s, t) - c(t - \gamma s)) > u \right), \end{aligned}$$

where $Z(s, t) := X_H(t) - \gamma X_H(s)$, $s, t \geq 0$. Using the self-similarity of the fBm X_H , we further have

$$\psi_{\gamma, T_u}(u) = \mathbb{P} \left(\sup_{0 \leq s \leq t \leq 1} Y_u(s, t) > \frac{u}{T_u^H} \right), \quad (12)$$

where, for any $u > 0$

$$Y_u(s, t) = \frac{Z(s, t)}{1 + \frac{cT_u}{u}(t - \gamma s)}, \quad s, t \geq 0. \quad (13)$$

In order to prove statement i) in Theorem 2.2, we give the following crucial lemma.

Lemma 3.1 Let $\{Y_u(s, t), s, t \geq 0\}, u > 0$ be a family of Gaussian random fields defined as in (13) with $H \in (0, 1)$ and $\gamma \in (0, 1)$. Assume that the condition of statement i) in Theorem 2.2

is satisfied. Then, for any u large enough, the variance function $V_{Y_u}^2(s, t) = \mathbb{E}(Y_u^2(s, t))$ of the Gaussian random field Y_u attains its maximum over the set $A := \{(s, t) : 0 \leq s \leq t \leq 1\}$ at the unique point $(0, 1)$. Moreover,

$$V_{Y_u}(0, 1) = \frac{u}{u + cT_u}.$$

Proof of Lemma 3.1 We only present the main ideas of the proof omitting thus some tedious and straightforward calculations. By solving the two equations

$$\frac{\partial V_{Y_u}^2(s, t)}{\partial s} = 0, \quad \frac{\partial V_{Y_u}^2(s, t)}{\partial t} = 0$$

we have that $s = t$. Therefore, the maximum of $V_{Y_u}^2(s, t)$ over A must be attained on the following three lines $l_1 = \{(0, t), 0 \leq t \leq 1\}$, $l_2 = \{(s, t), 0 \leq s = t \leq 1\}$ or $l_3 = \{(s, 1), 0 \leq s \leq 1\}$. It can be shown that on l_1 the maximum is attained uniquely at $(0, 1)$ and on l_2 the maximum is attained uniquely at $(1, 1)$. Obviously, both of the two points are on the line l_3 . Consequently, the maximum point of $V_{Y_u}^2(s, t)$ over A must be on l_3 . Moreover, we have that

$$(V_{Y_u}^2(s, 1))' = \frac{2c\gamma T_u}{u} \left(1 + \frac{cT_u}{u}(1 - \gamma s)\right)^{-3} f_{\frac{cT_u}{u}}(s),$$

where, for any $d > 0$

$$\begin{aligned} f_d(s) &= 1 - \gamma - (\gamma - \gamma^2)s^{2H} + \gamma(1 - s)^{2H} - \frac{H}{d}(1 + d - d\gamma s) \\ &\quad \times ((1 - \gamma)s^{2H-1} + (1 - s)^{2H-1}), \quad s \geq 0. \end{aligned} \tag{14}$$

Thus from the following technical lemma and the fact that

$$\lim_{u \rightarrow \infty} \frac{cT_u}{u} = cs_0 < \frac{H}{1 - H}$$

we conclude that

$$(V_{Y_u}^2(s, 1))' < 0, \quad \forall s \in (0, 1)$$

implying that the maximum of $V_{Y_u}^2(s, t)$ over the set A is attained at the unique point $(0, 1)$. This completes the proof. $\square$

Lemma 3.2 Let $f_d(s), s \geq 0$ be given as in (14) with $\gamma \in [0, 1)$ and $d \in [0, \frac{H}{1-H})$. Then

$$f_d(s) < 0, \quad \forall s \in (0, 1).$$

Proof of Lemma 3.2 First rewrite $f_d(s)$ as

$$f_d(s) = (1 - \gamma) + \gamma(1 - H)(1 - s)^{2H} - \gamma(1 - \gamma)(1 - H)s^{2H}$$

$$-H \left(1 + \frac{1}{d} - \gamma\right) (1-s)^{2H-1} - \frac{H}{d} (1+d)(1-\gamma)s^{2H-1}.$$

Further, we have

$$1 - \gamma < (1 - \gamma)(1 - s)^{2H-1} + (1 - \gamma)s^{2H-1}, \quad s \in (0, 1).$$

Then, replacing $1 - \gamma$ by $(1 - \gamma)(1 - s)^{2H-1} + (1 - \gamma)s^{2H-1}$ in the above equation we obtain

$$\begin{aligned} f_d(s) &< (1 - \gamma)(1 - s)^{2H-1} + (1 - \gamma)s^{2H-1} + \gamma(1 - H)(1 - s)^{2H} - \gamma(1 - \gamma)(1 - H)s^{2H} \\ &\quad - H \left(1 + \frac{1}{d} - \gamma\right) (1-s)^{2H-1} - \frac{H}{d} (1+d)(1-\gamma)s^{2H-1} \\ &< \left(1 - H - \frac{H}{d}\right) ((1-s)^{2H-1} + (1-\gamma)s^{2H-1}) - \gamma(1-\gamma)(1-H)s^{2H}, \end{aligned}$$

where in the second inequality we used the fact that

$$\gamma(1 - H)(1 - s)^{2H} \leq \gamma(1 - H)(1 - s)^{2H-1}, \quad \forall s \in (0, 1).$$

Since for any $d \in [0, \frac{H}{1-H})$

$$1 - H < \frac{H}{d}$$

we conclude that $f_d(s) < 0$ holds for all $s \in (0, 1)$, establishing the proof. $\square$

Proof of Theorem 2.2 *i*). First, note that (12) can be rewritten as

$$\psi_{\gamma, T_u}(u) = \mathbb{P} \left(\sup_{0 \leq s \leq t \leq 1} \frac{Y_u(s, t)}{V_{Y_u}(0, 1)} > \frac{u + cT_u}{T_u^H} \right).$$

Next, for any fixed large u , we give expansion of $\frac{V_{Y_u}(s, t)}{V_{Y_u}(0, 1)}$ at the point $(0, 1)$. It follows that

$$\frac{V_{Y_u}(s, t)}{V_{Y_u}(0, 1)} = \begin{cases} 1 - (H - c(u))(1 - t) - \gamma(H - c(u))s + o(1 - t + s), & H > 1/2, \\ 1 - (\frac{1}{2} - c(u))(1 - t) - \gamma(1 - \frac{\gamma}{2} - c(u))s + o(1 - t + s), & H = 1/2, \\ 1 - (H - c(u))(1 - t) - \frac{\gamma - \gamma^2}{2} s^{2H} + o(1 - t + s^{2H}), & H < 1/2 \end{cases} \quad (15)$$

holds as $(s, t) \rightarrow (0, 1)$, where $c(u) = \frac{cT_u}{u + cT_u}$. Furthermore, we have that

$$1 - Cov \left(\frac{Y_u(s, t)}{V_{Y_u}(s, t)}, \frac{Y_u(s', t')}{V_{Y_u}(s', t')} \right) = \frac{1}{2} (|t - t'|^{2H} + \gamma^2 |s - s'|^{2H}) (1 + o(1)) \quad (16)$$

holds as $(s, t), (s', t') \rightarrow (0, 1)$. In addition, there exists a positive constant $\mathbb{Q}$ such that, for all u large enough

$$\mathbb{E} \left(\left(\frac{Y_u(s, t)}{V_{Y_u}(0, 1)} - \frac{Y_u(s', t')}{V_{Y_u}(0, 1)} \right)^2 \right) \leq \mathbb{Q} (|t - t'|^{2H} + |s - s'|^{2H})$$

holds for all $(s, t) \in A$. Therefore, by the fact that

$$\lim_{u \rightarrow \infty} c(u) = c_0 = \frac{cs_0}{1 + cs_0} < H$$

and using Theorem 4.1 (see Appendix), we obtain that

$$\psi_{\gamma, T_u}(u) = D_{H, \gamma} \left(\frac{u + cT_u}{T_u^H} \right)^{\left(\frac{1-2H}{H}\right)+} \Psi \left(\frac{u + cT_u}{T_u^H} \right) (1 + o(1)) \quad (17)$$

as $u \rightarrow \infty$, where

$$\mathcal{D}_{H, \gamma} = \begin{cases} 2^{-\frac{1}{2H}} (H - c_0)^{-1} \mathcal{H}_{2H} \mathcal{P}_{2H}^{\frac{1-\gamma}{2H}}, & \text{if } H < 1/2, \\ \frac{4(1-c_0)^2}{(1-2c_0)(2-2c_0-\gamma)}, & \text{if } H = 1/2, \\ 1 & \text{if } H > 1/2. \end{cases}$$

Combining the above formula with (6) we obtain (9).

Next, we present the proof of statement *ii*).

Assume first that $\lim_{u \rightarrow \infty} \frac{T_u - t_0 u}{A(u)} = x \in \mathbb{R}$. We have from (4) that

$$\mathbb{P} \left(\frac{\tau_{\gamma, u} - t_0 u}{A(u)} \leq x \mid \tau_{\gamma, u} < \infty \right) \rightarrow \Phi(x)$$

holds as $u \rightarrow \infty$. Further note that the above is equivalent to

$$\lim_{u \rightarrow \infty} \frac{\mathbb{P} \left(\sup_{0 \leq t \leq t_0 u + x A(u)} W_{\gamma}(t) > u \right)}{\mathbb{P}(\tau_{\gamma, u} < \infty)} = \Phi(x).$$

Thus, we obtain that

$$\psi_{\gamma, T_u}(u) = \psi_{\gamma, \infty}(u) \Phi(x) (1 + o(1))$$

as $u \rightarrow \infty$, which together with (7) and Theorem 1.1 in [19] yields the validity of (10). Finally,

assume that $\lim_{u \rightarrow \infty} \frac{T_u - t_0 u}{A(u)} = \infty$. For any positive large M

$$\psi_{\gamma, t_0 u + M A(u)}(u) \leq \psi_{\gamma, T_u}(u) \leq \psi_{\gamma, \infty}(u)$$

holds for all u large enough, hence

$$\Phi(M) \leq \liminf_{u \rightarrow \infty} \frac{\psi_{\gamma, T_u}(u)}{\psi_{\gamma, \infty}(u)} \leq \limsup_{u \rightarrow \infty} \frac{\psi_{\gamma, T_u}(u)}{\psi_{\gamma, \infty}(u)} \leq 1.$$

Letting $M \rightarrow \infty$ in the above we conclude that

$$\psi_{\gamma, T_u}(u) = \psi_{\gamma, \infty}(u) (1 + o(1))$$

holds as $u \rightarrow \infty$, which further implies that (10) is valid, establishing the proof. $\square$

Proof of Theorem 2.3 We start with the proof of statement *i*). It follows from (17) that, for

any $x > 0$

$$\mathbb{P} \left(\frac{u^2 (T_u - \tau_{\gamma, u})}{T_u^{2H+1}} > x \mid \tau_{\gamma, u} \leq T_u \right) = \frac{\mathbb{P} \left(\sup_{0 \leq t \leq T_x(u)} W_{\gamma}(t) > u \right)}{\mathbb{P} \left(\sup_{0 \leq t \leq T_u} W_{\gamma}(t) > u \right)}$$

$$\begin{aligned}
&= \frac{D_{H,\gamma} \left(\frac{u+cT_x(u)}{(T_x(u))^H} \right)^{\left(\frac{1-2H}{H}\right)^+} \Psi \left(\frac{u+cT_x(u)}{(T_x(u))^H} \right)}{D_{H,\gamma} \left(\frac{u+cT_u}{T_u^H} \right)^{\left(\frac{1-2H}{H}\right)^+} \Psi \left(\frac{u+cT_u}{T_u^H} \right)} (1+o(1)) \\
&= \exp \left(-\frac{\left(\frac{u+cT_x(u)}{(T_x(u))^H} \right)^2 - \left(\frac{u+cT_u}{T_u^H} \right)^2}{2} \right) (1+o(1)) \\
&\rightarrow \exp(-Hx)
\end{aligned}$$

holds as $u \rightarrow \infty$, where $T_x(u) = T_u - xT_u^{2H+1}/u^2$. Therefore the claim follows.

Next, we give the proof of statement *ii*). Similar arguments as above yield that, for any $x > 0$

$$\begin{aligned}
\mathbb{P} \left(\frac{T_u - \tau_{\gamma,u}}{u^{2H-1}} > x \mid \tau_{\gamma,u} \leq T_u \right) &= \exp \left(-\frac{\left(\frac{u+c(T_u-xu^{2H-1})}{(T_u-xu^{2H-1})^H} \right)^2 - \left(\frac{u+cT_u}{T_u^H} \right)^2}{2} \right) (1+o(1)) \\
&\rightarrow \exp(-\lambda x)
\end{aligned}$$

holds as $u \rightarrow \infty$, where $\lambda = \frac{(1+cs_0)(H-(1-H)cs_0)}{s_0^{2H+1}}$. Finally, since by (10) for any $y \leq x$

$$\begin{aligned}
\mathbb{P} \left(\frac{\tau_{\gamma,u} - t_0 u}{A(u)} < y \mid \tau_{\gamma,u} \leq T_u \right) &= \frac{\mathbb{P} \left(\sup_{0 \leq t \leq t_0 u + yA(u)} W_\gamma(t) > u \right)}{\mathbb{P} \left(\sup_{0 \leq t \leq T_u} W_\gamma(t) > u \right)} \\
&\rightarrow \frac{\Phi(y)}{\Phi(x)}
\end{aligned}$$

holds as $u \rightarrow \infty$, the claim of statement *iii*) follows, and thus the proof is complete. $\square$

4 Appendix: Piterbarg's Theorem for Non-homogeneous Gaussian Fields

We present below a generalization of Theorem D.3 and Theorem 8.2 in [24], which is tailored for the proof of the main results. We first introduce a generalization of the Piterbarg constant given by

$$\tilde{\mathcal{P}}_\alpha^b = \lim_{S \rightarrow \infty} \mathbb{E} \left(\exp \left(\sup_{t \in [-S, S]} \left(\sqrt{2} B_\alpha(t) - (1+b)|t|^\alpha \right) \right) \right) \in (0, \infty), \quad \alpha \in (0, 2], \quad b > 0,$$

where $\{B_\alpha(t), t \in \mathbb{R}\}$ is a standard fBm defined on $\mathbb{R}$. Let $D = \{(s, t), 0 \leq s \leq t \leq 1\}$, and let $\{\eta_u(s, t), (s, t) \in D\}, u \geq 0$ be a family of Gaussian random fields satisfying the following three assumptions:

A1: The variance function $\sigma_{\eta_u}^2(s, t)$ of η_u attains its maximum on the set D at some unique point (s_0, t_0) for any u large enough, and further there exist four positive constants $A_i, \beta_i, i = 1, 2$ and

two functions $A_i(u), i = 1, 2$ satisfying $\lim_{u \rightarrow \infty} A_i(u) = A_i, i = 1, 2$ such that $\sigma_{\eta_u}(s, t)$ has the following expansion around (s_0, t_0) for all u large enough

$$\sigma_{\eta_u}(s, t) = 1 - A_1(u)|s - s_0|^{\beta_1}(1 + o(1)) - A_2(u)|t - t_0|^{\beta_2}(1 + o(1)), \quad (s, t) \rightarrow (s_0, t_0).$$

A2: There exist four constants $B_i > 0, \alpha_i \in (0, 2], i = 1, 2$ and two functions $B_i(u), i = 1, 2$ satisfying $\lim_{u \rightarrow \infty} B_i(u) = B_i, i = 1, 2$ such that the correlation function $r_{\eta_u}(s, t; s', t')$ of η_u has the following expansion around (s_0, t_0) for all u large enough

$$r_{\eta_u}(s, t; s', t') = 1 - B_1(u)|s - s'|^{\alpha_1}(1 + o(1)) - B_2(u)|s - s'|^{\alpha_2}(1 + o(1)), \quad (s, t), (s', t') \rightarrow (s_0, t_0).$$

A3: For some positive constants $\mathbb{Q}$ and γ , and all u large enough

$$\mathbb{E}(\eta_u(s, t) - \eta_u(s', t'))^2 \leq \mathbb{Q}(|s - s'|^\gamma + |t - t'|^\gamma)$$

for any $(s, t), (s', t') \in D$.

Theorem 4.1 *If $\{\eta_u(s, t), (s, t) \in D\}$, $u \geq 0$ is a family of Gaussian random fields satisfying **A1-A3**, then*

$$\mathbb{P}\left(\sup_{(s, t) \in D} \eta_u(s, t) > u\right) = \mathcal{F}_{\alpha, \beta}^{(1)}(u) \mathcal{F}_{\alpha, \beta}^{(2)}(u) \Psi(u), \quad \text{as } u \rightarrow \infty,$$

where

$$\mathcal{F}_{\alpha, \beta}^{(i)}(u) = \begin{cases} \widehat{I}_i \mathcal{H}_{\alpha_i} B_i^{\frac{1}{\alpha_i}} A_i^{-\frac{1}{\beta_i}} \Gamma\left(\frac{1}{\beta_i} + 1\right) u^{\frac{2}{\alpha_i} - \frac{2}{\beta_i}}, & \text{if } \alpha_i < \beta_i, \\ \widehat{\mathcal{P}}_{\alpha_i}^{\frac{A_i}{B_i}}, & \text{if } \alpha_i = \beta_i, \\ 1 & \text{if } \alpha_i > \beta_i, \end{cases} \quad i = 1, 2,$$

with $\Gamma(\cdot)$ the Euler Gamma function and

$$\widehat{\mathcal{P}}_{\alpha_1}^{\frac{A_1}{B_1}} = \begin{cases} \widehat{\mathcal{P}}_{\alpha_1}^{\frac{A_1}{B_1}}, & \text{if } s_0 \in (0, 1), \\ \mathcal{P}_{\alpha_1}^{\frac{A_1}{B_1}} & \text{if } s_0 = 0 \text{ or } 1, \end{cases} \quad \widehat{\mathcal{P}}_{\alpha_2}^{\frac{A_2}{B_2}} = \begin{cases} \widehat{\mathcal{P}}_{\alpha_2}^{\frac{A_2}{B_2}}, & \text{if } t_0 \in (0, 1), \\ \mathcal{P}_{\alpha_2}^{\frac{A_2}{B_2}} & \text{if } t_0 = 0 \text{ or } 1, \end{cases}$$

$$\widehat{I}_1 = \begin{cases} 2, & \text{if } s_0 \in (0, 1), \\ 1 & \text{if } s_0 = 0 \text{ or } 1, \end{cases} \quad \widehat{I}_2 = \begin{cases} 2, & \text{if } t_0 \in (0, 1), \\ 1 & \text{if } t_0 = 0 \text{ or } 1. \end{cases}$$

Proof of Theorem 4.1 It follows from the assumptions **A1-A2** that for any $\varepsilon > 0$ and for u large enough we have

$$(A_1 - \varepsilon)|s - s_0|^{\beta_1} + (A_2 - \varepsilon)|t - t_0|^{\beta_2} \leq 1 - \sigma_{\eta_u}(s, t) \leq (A_1 + \varepsilon)|s - s_0|^{\beta_1} + (A_2 + \varepsilon)|t - t_0|^{\beta_2}$$

as $(s, t) \rightarrow (s_0, t_0)$, and

$$(B_1 - \varepsilon)|s - s'|^{\alpha_1} + (B_2 - \varepsilon)|t - t'|^{\alpha_2} \leq 1 - r_{\eta_u}(s, t; s', t') \leq (B_1 + \varepsilon)|s - s'|^{\alpha_1} + (B_2 + \varepsilon)|t - t'|^{\alpha_2}$$

as $(s, t), (s', t') \rightarrow (s_0, t_0)$. Therefore, in the light of Theorem 8.2 in [24] we can get asymptotical upper and lower bounds, and thus the claims follow by letting $\varepsilon \rightarrow 0$. The proof is complete. $\square$

Acknowledgement: The authors kindly acknowledge partial support from the Swiss National Science Foundation Project 200021-140633/1, and the project RARE -318984 (an FP7 Marie Curie IRSES Fellowship).

References

- [1] J.M.P. Albin and H. Choi. A new proof of an old result by Pickands. *Electronic Communications in Probability*, 15:339–345, 2010.
- [2] S. Asmussen. *Applied probability and queues*. Wiley Series in Probability and Mathematical Statistics: Applied Probability and Statistics. John Wiley & Sons Ltd., Chichester, 1987.
- [3] S. Asmussen and H. Albrecher. *Ruin probabilities*. Advanced Series on Statistical Science & Applied Probability, 14. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, second edition, 2010.
- [4] H. Awad and P. Glynn. Conditional limit theorems for regulated fractional Brownian motion. *Ann. Appl. Probab.*, 19(6):2102–2136, 2009.
- [5] P. Boulongne, D. Pierre-Loti-Viaud, and V.I. Piterbarg. On average losses in the ruin problem with fractional Brownian motion as input. *Extremes*, 12(1):77–91, 2009.
- [6] K. Dębicki and P. Kisowski. A note on upper estimates for Pickands constants. *Statistics & Probability Letters*, 78(14):2046–2051, 2008.
- [7] K. Dębicki. Ruin probability for Gaussian integrated processes. *Stochastic Process. Appl.*, 98(1):151–174, 2002.
- [8] K. Dębicki, E. Hashorva, and L. Ji. Gaussian risk model with financial constraints. *Scandinavian Actuarial Journal*, in press, 2014.
- [9] K. Dębicki and K. Kosiński. On the infimum attained by the reflected fractional Brownian motion. *Preprint*, 2014.
- [10] K. Dębicki and M. Mandjes. Exact overflow asymptotics for queues with many Gaussian inputs. *J. Appl. Probab.*, 40(3):704–720, 2003.

- [11] K. Dębicki and T. Rolski. A note on transient Gaussian fluid models. *Queueing Systems, Theory and Applications*, 42:321–342, 2002.
- [12] A.B. Dieker. Extremes of Gaussian processes over an infinite horizon. *Stochastic Process. Appl.*, 115(2):207–248, 2005.
- [13] A.B. Dieker and B. Yakir. On asymptotic constants in the theory of Gaussian processes. *Bernoulli*, to appear, 2014.
- [14] P. Embrechts, C. Klüppelberg, and T. Mikosch. *Modelling extremal events*, volume 33 of *Applications of Mathematics (New York)*. Springer-Verlag, Berlin, 1997. For insurance and finance.
- [15] P.S. Griffin. Convolution equivalent Lévy processes and first passage times. *Ann. Appl. Probab.*, 23:1506–1543, 2013.
- [16] P.S. Griffin and R.A. Maller. Path decomposition of ruinous behaviour for a general Lévy insurance risk process. *Ann. Appl. Probab.*, 22:1411–1449, 2012.
- [17] J. M. Harrison. *Brownian motion and stochastic flow systems*. Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics. John Wiley & Sons Inc., New York, 1985.
- [18] E. Hashorva and L. Ji. Approximation of passage times of γ -reflected processes with fBm input. *J. Appl. Probab.*, to appear, 2014.
- [19] E. Hashorva, L. Ji, and V.I. Piterbarg. On the supremum of γ -reflected processes with fractional Brownian motion as input. *Stochastic Process. Appl.*, 123:4111–4127, 2013.
- [20] J. Hüsler and V.I. Piterbarg. Extremes of a certain class of Gaussian processes. *Stochastic Process. Appl.*, 83(2):257–271, 1999.
- [21] J. Hüsler and V.I. Piterbarg. A limit theorem for the time of ruin in a Gaussian ruin problem. *Stochastic Process. Appl.*, 118(11):2014–2021, 2008.
- [22] J. Hüsler and Y. Zhang. On first and last ruin times of Gaussian processes. *Statistics & Probability Letters*, 78(10):1230–1235, 2008.

- [23] S. G. Kobelkov. A limit theorem for the ruin time with power profits in the case of an integrated Gaussian stationary process. *Vestnik Moskov. Univ. Ser. I Mat. Mekh.*, (4):3–11, 2011.
- [24] V.I. Piterbarg. *Asymptotic methods in the theory of Gaussian processes and fields*, volume 148 of *Translations of Mathematical Monographs*. American Mathematical Society, Providence, RI, 1996.
- [25] W. Whitt. *Stochastic-process limits*. Springer Series in Operations Research. Springer-Verlag, New York, 2002.
- [26] A. J. Zeevi and P. W. Glynn. On the maximum workload of a queue fed by fractional Brownian motion. *Ann. Appl. Probab.*, 10(4):1084–1099, 2000.